National Childhood Immunisation Schedule (NCIS)

(from birth to age 17 years, effective from 1 November 2020)

Venetee	Di d				40	45	10	2.4		10.11	42.42	12.11	45.47
Vaccine	Birth	2 months	4 Months	6 months	12 months	15 months	18 months	2-4	5-9	10-11	12-13	13-14	15-17
		months	iviontns	months	months	months	months	years	years	years	years	years	years
Bacillus Calmette-Guérin (BCG)	D1												
Hepatitis B (HepB)	D1	D2		D3									
Diphtheria, tetanus and acellular		D1	D2	D3			B1						
pertussis (paediatric) (DTaP)													
Tetanus, reduced diphtheria and										B2			
acellular pertussis (Tdap)													
Inactivated poliovirus (IPV)		D1	D2	D3			B1			B2			
mactivated pollovirus (ir v)		01	02	03			D1			02			
		54											
Haemophilus influenzae type b		D1	D2	D3			B1						
(Hib)													
Pneumococcal conjugate			D1	D2	B1								
(PCV10 or PCV13)													
Pneumococcal polysaccharide								One or two doses for children and adolescents age 2-17 years with specific					
(PPSV23)								medical condition or indication.					
Measles, mumps and rubella					D1	D2							
(MMR)													
Varicella (VAR)					D1	D2							
, ,													
Human papillomavirus											D1	D2	
(HPV2 or HPV4)											(Females)	(Females)	
Influenza (INF)				Annual vaccination or per season for all children age 6 months to									
IIIIIdeliza (IIVF)				<5 years (6-59 months).					age 5-17 years with specific medical condition or indication.				
				<5 years (0-59 months).					age 3-17 years with specific medical condition of indication.				



Recommended ages and doses for all children



Recommended for persons with specific medical condition or indication

FOOTNOTES:

- D1, D2, D3: Dose 1, dose 2, dose 3
- B1, B2: Booster 1, booster 2
- 10-11, 12-13, 13-14 years: Primary 5, Secondary 1, Secondary 2 (Tdap, IPV, HPV (for females) and MMR (as catch-up) vaccines are provided as part of Health Promotion Board's school-based vaccination programme)
- HepB: Doses 2 and 3 are recommended to be given as part of the 6-in-1 vaccine at 2 and 6 months, respectively
- MMR: Only the dose 2 is recommended to be given as part of the MMRV vaccine